

Postnatal Bladder Care Guideline

Document Type:	Clinical Guidelines
Ref:	(For Non-Clinical References – Contact: CTM Corporate Governance@wales.nhs.uk For Clinical References – Contact: CTM ClinicalPolicies@wales.nhs.uk
Author:	Nadia Hikary Bhal, Rajul Patel- Gardia
Executive Sponsor:	Choose an item.
Approved By:	Choose an item.
Approval / Effective Date:	October 2024
Review Date:	October 2027
Version:	2

Target Audience:

People who need to know about this document in detail	All Midwifery, obstetric and gynaecology staff working within CTM UHB
People who need to have a broad understanding of this document	As above
People who need to know that this document exists	As above

Integrated Impact Assessment:

Equality Impact Assessment Date & Outcome	Date: November 2024
	Outcome: there is no negative impact
Welsh Language Standard	Choose an item.
Date of approval by Equality Team:	(00/00/0000)
Aligns to the following Wellbeing of Future Generation Act Objective	Choose an item.



Disclaimer:

If the review date of this document has passed please ensure that the version you are using is the most up to date version by checking on WISDOM

Guidelines Definition

Clinical guidelines are systemically developed statements that assist clinicians and patients in making decisions about appropriate treatments for specific conditions.

They allow deviation from a prescribed pathway according to the individual circumstances and where reasons can be clearly demonstrated and documented.

Minor Amendments

If a minor change is required to the document, which does not require a full review please identify the change below and update the version number.

Type of change	Why change made	Page number	Date of change	Version 1 to 1.1	Name of responsible person
Update /review of guideline					Nadia Hikary Bhal, Rajul Patel- Gardia

Equality Impact Assessment Statement

This Procedure has been subject to a full equality assessment and no impact has been identified.

Contents

Introduction and Aim	4
Objectives	4
Scope	4
Risk Factors for Postpartum Voiding Dysfunction	Error! Bookmark not defined.
Intrapartum bladder care	5
Postpartum bladder care	6
Symptoms of Postpartum Voiding Dysfunction.....	7
Hospital Birth	7
Home Birth	9
CTMUHB Postpartum Bladder Care Pathway.....	10
CTMUHB Referral form	11
References	12

Introduction and Aim

Hormonal changes in pregnancy decrease the tone of the detrusor muscle. Combined with trauma to the bladder, pelvic floor muscles and nerves during delivery the postpartum bladder tends to become underactive and is therefore vulnerable to the retention of urine.

Postpartum voiding dysfunction is defined as failure to pass urine spontaneously within 6 hours of vaginal delivery or following catheter removal after delivery. This occurs in 0.7-4% of deliveries.

Overt urinary retention is the inability to void postpartum.

Covert urinary retention occurs when there is an elevated post-void residual volume (PVRV) of >150mL urine *without* symptoms of urinary retention.

If postpartum voiding dysfunction is unrecognised it can lead to long term sequelae such as recurrent urinary tract infection and urinary incontinence. Prevention of this should aim to identify all women unable to pass urine within 6 hours of delivery or catheter removal *and* all women who are symptomatic of voiding dysfunction.

The importance of prompt diagnosis and appropriate management of these women cannot be over-emphasised as early intervention is the key to ensuring rapid return to normal bladder function.

Objectives

To maintain bladder function and to provide appropriate management to women with postpartum voiding dysfunction.

To minimise the risk of prolonged voiding dysfunction in the puerperium and prevent its long term sequelae.

Scope

This policy applies to all clinicians working within maternity services including temporary staff, locums, bank and agency / annualised hours staff and visiting clinicians.

Risk Factors for Postpartum voiding dysfunction

Many risk factors have been identified for the development of postpartum voiding dysfunction, including the following:

- Primiparity
- Assisted birth
- Epidural analgesia
- Prolonged labour
- Perineal trauma
- History of voiding problems

Postpartum voiding dysfunction however can develop in women without any identifiable risk factors, regardless of mode of birth or analgesia.

Intrapartum bladder care

Prevention of postpartum voiding dysfunction starts with good bladder management during intrapartum care. This includes the documentation of frequency and volume of bladder emptying. Women should be encouraged to empty their bladder at regular intervals. NICE (2023) advises bladder care should be reviewed at least every 4 hours and should include:-

- Frequency of passing urine and bladder sensation.
- Fluid balance monitoring if sensation is abnormal or absent. If there is an inability to pass urine, or the woman is receiving intravenous fluids.
- Offering to insert a catheter if there are any ongoing concerns over the woman's ability to pass urine.

If an intermittent catheter is to be used this ideally is done at the time of vaginal examination. If catheterisation is likely to be used more than twice during labour, an indwelling catheter should be considered. An indwelling catheter should be removed during the active second stage of labour and assisted vaginal birth to prevent trauma to the bladder and urethra.

Postpartum bladder care

Consider (re-)inserting an in-dwelling urinary catheter in women after:

- Regional anaesthesia and prolonged labour
- Mid-cavity assisted birth
- Urethral trauma
- Severe perineal trauma
- Women receiving High Dependency Care
- For all births and procedures in theatre, who have spinal anaesthesia (including combined spinal-epidural) or who have had epidural anaesthesia “topped up”.

Insertion of a catheter should be documented on the local Urinary Catheter Bundle pathway.

Regional anaesthesia can affect bladder sensation and therefore the indwelling catheter should not be removed until the woman is mobile as a minimum unless specified otherwise in the operation note. It may be appropriate to leave an indwelling catheter in place for a longer period for example if there is significant perineal trauma/oedema or there is a need for accurate measurement of the urine output e.g. intra or post postpartum sepsis

All women should void within 6 hours of birth or indwelling catheter removal.

The Royal College of Obstetricians and Gynaecologists (RCOG) study group on incontinence recommends that no woman should be allowed to go longer than 6 hours without voiding or catheterisation postpartum. Encouragement to pass urine after 4 hours allows time for conservative measures (analgesia, mobilisation, bath or shower, privacy).

Symptoms of Postpartum Voiding Dysfunction

Signs and symptoms that should raise the alert to voiding dysfunction include:

- Inability to pass urine within 6 hours of birth or catheter removal
- Slow urinary stream
- Urinary frequency passing small amounts of urine
- (Sensation of) incomplete emptying
- Urinary incontinence

It is important to recognise that acute retention can be **painless** in the postpartum period especially following regional analgesia.

Hospital Birth

1. Document time and volume of first void after birth or after removal of indwelling catheter on the postnatal pathway. It is important to ask the woman about her voiding pattern as this could point towards voiding dysfunction (see symptoms above). In women with an indwelling catheter time of removal of the catheter must also be documented in the postnatal pathway and urinary catheter bundle.

2. Insert an in/out catheter OR bladder scan for post void residual volume (PVRV) in the following women:

- No void within 6 hours of birth or removal of indwelling catheter. OR
- Passing frequent small amounts of urine with the sensation of incomplete voiding.

A bladder scan however, may not give accurate readings in women with a high BMI or with the presence of clots in the uterus.

2a. If PVRV < 150mL: no further management needed in the asymptomatic patient.

If PVRV < 500mL > 150: measure the next voided volume and PVRV.

2b. If PVRV > 500 mL or > 20% of voided volume: insert an indwelling catheter for 24 – 48 hours followed by trial without catheter (TWOC) – this can be done as an outpatient.

The obstetric team should be informed.

3. If at TWOC the woman is either unable to void within 6 hours or has a PVRV > 150mL; record the next 2 voids and if PVRV > 150mL after the 2nd void then re-catheterise the woman for 1 week.

Leave the catheter on free drainage. TWOC should be attempted after 1 week. (This can be done as an outpatient)

4. At 2nd TWOC record 2 voids and if the woman is either unable to void within six hours or has a PVRV > 150mL after 2nd void; re-catheterise designed for one week and complete a referral to **Sister Angharad Carroll** –local unit Urogynaecology Nurse Practitioner at PCH or **Sister Vickie Atkins** Urogynaecology nurse at POWH (see page 11 for referral form) who will liaise with the patient's local Urogynaecology consultant for further management plan.

Above management of postpartum retention and voiding dysfunction is summarised in attached flow diagram.

In all of these cases, the time and volume of voiding must be documented in the hospital notes. The voided volumes and the PVRV must also be recorded. Measurement of intake and output volumes needs to be recorded in these cases and a fluid balance chart commenced.

Further management aims to identify any factors contributing to delayed bladder emptying and to ensure adequate bladder drainage while waiting for normal function to return. **Following the diagnosis of urinary**

retention or voiding dysfunction, the following actions should be taken and documented in the hospital notes:

- Perform urinalysis and sent for MC&S as the presence of infection is an important contributory factor to prolonged voiding dysfunction.
- If a urinary tract infection is suspected, prompt antibiotic therapy should be initiated following Microguide guidance.
- The perineum should be examined and if swollen or painful, a catheter should be sited until the swelling and pain have settled.
- Ensure and provide adequate analgesia, as perineal pain is a significant factor in development of retention.
- Avoid and treat constipation if required.

All women experiencing voiding dysfunction must have follow up after discussion with the responsible consultant or senior registrar. It is the responsibility of the midwife who discharges the woman from the postnatal area to ensure that an appointment to see the UroGynaecology nurse practitioner is made as per the flow chart.


Home Birth

Following a homebirth, the woman should be instructed to make a note of the time of the first void and contact the community midwife if

- this has not occurred within 6 hours or
- there are any symptoms of voiding dysfunction

Referral to the postnatal ward can be arranged for management as described above for a hospital birth.

**CWM TAF MORGANNWG
UNIVERSITY HEALTH BOARD
POSTNATAL BLADDER CARE FLOWCHART**



**CWM TAF MORGANNWG UNIVERSITY HEALTH BOARD
POST NATAL VOIDING DYSFUNCTION
REFERRAL FORM**

Pt sticker

Referral date.....

Consultant.....

Patient's tel no.....

Patient BMI:

Date of delivery:

Parity:

Baby Weight :

Vaginal birth Forceps Ventouse Epidural Caesarean birth
3rd/4th Degree Tear

History of presenting complaint:

.....
.....

Date of initial catheterisation.....

Date of 1st TWOCRecatheterised

Date of 2nd TWOC Recatheterised Flip Flo valve attached: YES / NO

Date of 3rd TWOCRecatheterised Flip Flo Valve attached: YES /NO

(Flip Flow Valve during day time and Bladder can be left on free drainage at night)

Signature of Person Referring:

PRINT NAME & Designation:

Referrers Ext/Contact No.

**PLEASE COMPLETE THIS FORM AND ENSURE YOU HAVE THE PATIENTS
CORRECT CONTACT NUMBER. PLEASE FORWARD TO:**

**FAO – Sister Angharad Carroll – PCH
Sister Vickie Atkins - POW**

Incomplete or Illegible forms will be returned to sender

References

NICE guideline (NG235) Intrapartum care 2023

Kearney R, Cutner A (2008) Postpartum Voiding Dysfunction. *The Obstetrician & Gynaecologist* 10:2:71, 71-74

Ching-Chung et al (2002) Postpartum urinary retention: assessment of contributing factors and long term clinical impact. *Australian & New Zealand Journal of Obstetrics & Gynaecology* 42(4) 365-8

Dorflinger A & Monga A (2001) Voiding dysfunction. *Current Opinion in Obstetrics & Gynaecology* 13 507-512

Jeffery et al (1990) Chronic urinary retention postpartum. *Australian & New Zealand Journal of Obstetrics & Gynaecology*. Nov 30 (4) 364-366

Watson W J (1991) Prolonged postpartum urinary retention. *Military medicine* 156(9) 502-503

Carley et al (2002) Factors that are associated with clinically overt postpartum urinary retention after vaginal delivery. *American Journal of Obstetrics & Gynaecology* Aug 187(2) 430-433

Glavin K & Bjork J (2003) Incidence and treatment of urinary retention postpartum. *International Urogynaecology Journal* 14(2) 119-121

Zaki et al (2004) National survey for intrapartum and postpartum bladder care: assessing the need for guidelines. *BJOG Monitoring compliance and effectiveness*

Khullar V & Cardozo L D (2002) Bladder sensation after epidural analgesia. *Neurourology Urodynamics* 1993; 89: 424-425. MacLean A & Cardozo L. *Incontinence in Women*, London, RCOG Press

Ramsay I & Torbet T (1993) Incidence of abnormal voiding parameters in the immediate postpartum period. *Neurourology Urodynamics* 12: 179-183

Reo et al (2007) Clinically overt postpartum urinary retention after vaginal delivery: a retrospective case-control study. *International Urogynaecology Journal Pelvic Floor Dysfunction* 18: 521-524

