

Induction of Labour Guideline

Guideline information

Guideline number: 667

Classification:
Clinical

Supersedes:
Previous Versions

Local Safety Standard for Invasive Procedures (LOCSSIP) reference:
N/A

National Safety Standards for Invasive Procedures (NatSSIPs) standards:
N/A

Version number:
3.0

Date of Equality Impact Assessment:
Detail date of EqIA

Approval information

Approved by: Obstetric Written Documentation Review Group

Date of approval:
15/07/2022

Date made active:
03/01/2023

Review date:
15/07/2025

Summary of document:

This guideline is for use employees of Hywel Dda University Healthboard working in and alongside the Women Children's Health Directorate caring for all women and birthing people and birthing people who require induction of labour.

Scope:

This guideline is applicable to all women and birthing people where induction of labour is indicated when it is agreed that there is a higher probability of a healthier outcome for mother and birthing person and / or fetus to induce birth than if the pregnancy were to continue. Induction of labour should only be considered when vaginal birth is felt to be the most appropriate route.

The guidance uses the term “woman” (pronouns she or her) or Mother to describe individuals whose sex assigned at birth was female, whether they identify as female, male or non-binary. It is important to acknowledge it is not only people who identify as women for whom it is necessary to access women’s health and reproductive services. Therefore, this should include people who do not identify themselves as women but who are pregnant or have recently given birth. Obstetric and midwifery services and delivery of care must therefore be appropriate, inclusive and sensitive to the needs of those individuals whose gender identify does not align with the sex that they were assigned at birth.

To be read in conjunction with:

[623 - Large for Gestational Age in Non-Diabetic Guideline](#) (opens in new tab)

[621 - Hypertension in Pregnancy Guideline](#) (opens in new tab).

[669 - Management of the Small for Gestational Age Fetus Guideline](#) (opens in new tab).

[632 - Diabetes in Pregnancy Guideline](#) (opens in new tab)

Patient information:

[Include links to Patient Information Library](#)

Owning group:

Obstetric Written Documentation Review Group

15/07/2022

medicines formulary group signed off

Executive Director job title:

Mandy Rayani – Director of Nursing, Quality and Patient Experience

Reviews and updates:

1.0 – New Guideline

2.0 – Updated

3.0 – Full Review

Keywords

Induction of Labour, Postdates, Low Risk Pregnancy

Glossary of terms

OIL – Induction of Labour

VBAC - Vaginal Birth after Caesarean Section

SGA - Small for Gestational Age

IUGR - Intrauterine Growth Restriction

PPROM - Preterm Prelabour Rupture of Membranes

PROM - Prelabour Rupture of Membranes

ARM - Artificial Rupture of the Membranes

Contents

Guideline information	1
Approval information	1
Scope.....	5
Aim.....	5
Objectives	5
Introduction	5
Informing Decision Making	5
Explain to Women and Birthing People that there is a Small Increase in Risks of an Adverse Outcome associated with a Pregnancy continuing beyond 41+0 and these include:.....	6
Women and Birthing People over 40 Years of Age	6
Suspected Fetal Macrosomia (Large for Gestational Age)	6
History of Precipitate Labour	6
Maternal Request	7
Vaginal Birth after Caesarean Section (VBAC).....	7
Pre-Eclampsia /Hypertensive Disease in Pregnancy.....	7
Small for Gestational Age (SGA) / Intrauterine Growth Restriction (IUGR)	7
Preterm Prelabour Rupture of Membranes (PPROM) / Prelabour Rupture of Membranes (PROM) ...	7
Diabetes	7
Women and Birthing People who Decline Induction of Labour	7
Women and Birthing people who Decline Induction of Labour for Postmaturity (>41/40)	7
Methods of IOL	7
Membrane Sweeping.....	7
Cervical Ripening by Prostaglandins	8
Propess® should be Removed	8
Assessment	9
Process of Induction of Labour	9
Fetal Monitoring	10
Management of Uterine Hyperstimulation	11
Unsuccessful IOL.....	11
Further Cycle of IOL	11
Category 3 Caesarean Birth	12
Auditable Standards.....	12

References.....12

Appendix 1 – IOL Sticker14

Appendix 2 – Bishop’s Score Sticker15

Appendix 3 – Induction of Labour Information Leaflet (English).....16

Scope

This guideline is applicable to all women and birthing people where induction of labour is indicated when it is agreed that there is a higher probability of a healthier outcome for mother and birthing person and / or fetus to induce birth than if the pregnancy were to continue. Induction of labour should only be considered when vaginal birth is felt to be the most appropriate route.

The guidance uses the term “woman” (pronouns she or her) or Mother to describe individuals whose sex assigned at birth was female, whether they identify as female, male or non-binary. It is important to acknowledge it is not only people who identify as women for whom it is necessary to access women’s health and reproductive services. Therefore, this should include people who do not identify themselves as women but who are pregnant or have recently given birth. Obstetric and midwifery services and delivery of care must therefore be appropriate, inclusive and sensitive to the needs of those individuals whose gender identity does not align with the sex that they were assigned at birth.

Aim

The aim of this document is to:

- Define principles of induction of labour (IOL) for post maturity in low risk patients. IOL in special circumstances must be discussed with patient’s Consultant (Consultant on-call) or are discussed in specific guideline (IOL in the presence of uterine scar).

Objectives

The aim of this document will be achieved by the following objectives:

- Clearly defining IOL
- Detailing the methods of IOL
- Assessment of patients
- Process of induction

Introduction

Induction of Labour (IOL) is defined as an intervention designed to artificially initiate uterine contractions leading to progressive effacement and dilatation of the cervix, and birth of the baby. This may include women and birthing people and birthing people with membranes still intact or women and birthing people and birthing people with ruptured membranes who are not in labour.

Induction of labour is indicated when it is agreed that there is a higher probability of a healthier outcome for mother and birthing person and / or fetus to induce birth than if the pregnancy were to continue. Induction of labour should only be considered when vaginal birth is felt to be the most appropriate route.

Informing Decision Making

Before the induction process begins, women and birthing people should be informed about the clinical indication of the induction and the associated risks and benefits. Women and birthing people should also be informed of the arrangements for support and pain relief. Written information to support their decision making should be provided. Alternative options should a women and birthing people chose not to have an induction should be explained.

Women and birthing people should also be informed:

- Their hospital stay may be longer than with a spontaneous labour
- Vaginal examinations to assess the cervix are needed before and during induction
- Their choice of place of birth will be limited, as they may be recommended interventions (for example, oxytocin infusion, continuous fetal heart rate monitoring and epidurals) that are not available for home birth or in midwife-led birth units
- There may be limitations on the use of a birthing pool
- There may be a need for an assisted vaginal birth, with the associated increased risk of obstetric anal sphincter injury
- Pharmacological methods of induction can cause hyperstimulation
- An induced labour may be more painful than a spontaneous labour
- Induction may not be successful, and how this would affect the woman and birthing person's options

Explain to Women and Birthing People that there is a Small Increase in Risks of an Adverse Outcome associated with a Pregnancy continuing beyond 41+0 and these include:

- Increased likelihood of caesarean birth
- Increased likelihood of the baby needing admission to a neonatal intensive care unit
- Increased likelihood of stillbirth and neonatal death.

Therefore, in HDUHB induction of labour to be offered from 41/40

Decision to perform routine IOL after 41/40 for low risk women and birthing people (uncomplicated pregnancy, no concerning obstetric history) can be done by midwife or obstetrician and clear indication stated in notes and IOL file. In case of any complications in obstetric history this must be discussed with senior obstetrician.

IOL for any other reason must be discussed and agreed by named Consultant

Women and Birthing People over 40 Years of Age

Because there is some evidence that women and birthing people over 40 years of age have a slightly increased risk of intrauterine death, IOL can be considered after 39/40.

Women and birthing people over 40 years of age should have individualised plan for IOL made by their Consultant and documented in notes.

If a woman and birthing person over the age of 40 goes into spontaneous labour with no additional risk factors in pregnancy they can be treated as low risk in labour regardless of the birth setting. Note: Refer to All Wales Midwife Led Care Guideline.

Suspected Fetal Macrosomia (Large for Gestational Age)

In the absence of any other indications or problems (eg diabetes), suspected fetal macrosomia may be a consideration for induction of labour (Refer to [623 - Large for Gestational Age in Non-Diabetic Guideline](#) (opens in new tab) for further information).

History of Precipitate Labour

IOL to avoid a birth unattended by healthcare professionals should not be routinely offered to women and birthing people with a history of precipitate labour.

Maternal Request

Consider requests for induction of labour only after discussing the benefits and risks with the woman and birthing person, taking into account the woman and birthing person's circumstances and preferences.

Vaginal Birth after Caesarean Section (VBAC)

IOL after caesarean section carries the potential additional risk of uterine scar dehiscence/rupture. This must be agreed by the consultant obstetrician. See VBAC guideline.

Pre-Eclampsia /Hypertensive Disease in Pregnancy

Refer to [621 - Hypertension in Pregnancy Guideline](#) (opens in new tab).

Small for Gestational Age (SGA) / Intrauterine Growth Restriction (IUGR)

Refer to [669 - Management of the Small for Gestational Age Fetus Guideline](#) (opens in new tab).

Preterm Prelabour Rupture of Membranes (PPROM) / Prelabour Rupture of Membranes (PROM)

Refer to Preterm Prelabour Rupture of Membranes (PPROM) / PROM Prelabour Rupture Of Membranes guideline

Diabetes

Refer to [632 - Diabetes in Pregnancy Guideline](#) (opens in new tab)

Women and Birthing People who Decline Induction of Labour

Women and birthing people who decline IOL should be referred to a Consultant Obstetrician for individualised care planning

A clear management plan should be documented in the woman and birthing person's hand-held and maternity notes.

Women and Birthing people who Decline Induction of Labour for Postmaturity (>41/40)

From >42/40 women and birthing people should be offered increased antenatal monitoring consisting of computerised CTG (Dawes Redman) and ultrasound estimation of maximum amniotic pool depth twice weekly. Women and birthing people must be aware that CTG and USS surveillance has a poor predictive value.

Any abnormalities found on the CTG or liquor volume should be discussed with the woman and birthing person and the consultant on-call and recommendation for IOL should be discussed again.

Methods of IOL

The methods of induction are varied and success depends on appropriate assessment and treatment.

Membrane Sweeping

Explain to women and birthing people:

- What a membrane sweep is
- That membrane sweeping might make it more likely that labour will start without the need for additional pharmacological or mechanical methods of induction

- That pain, discomfort and vaginal bleeding are possible from the procedure.

At antenatal visits after 39+0 weeks, discuss with women and birthing people if they would like a vaginal examination for membrane sweeping, and if so obtain verbal consent from them before carrying out the membrane sweep.

Discuss with women and birthing people whether they would like to have additional membrane sweeping if labour does not start spontaneously following the first sweep (Refer to standard operating procedures for membrane sweeping in pregnancy).

Women and birthing people should be offered membrane sweeps in the week leading to induction of labour, this should not be performed prior to 37/40 – timing and frequency of sweeps should be discussed between the woman and birthing person and her midwife.

Cervical Ripening by Prostaglandins

Prostaglandins should be used to ripen the cervix prior to artificial rupture of the membranes if the woman and birthing person's Bishop score is 6 or less.

Prostin E2® Vaginal Gel (Dinoprostone)

Dinoprostone in the form of vaginal gel 1mg or 2mg should be considered as the preferred form in women and birthing people with a favourable Bishop score of between 4 and 6.

Standard regime considers administration of 1-2mg followed by a second dose of 1-2mg in 6-12 hours interval up to 4mg total dose in nulliparous women and birthing people and 3mg total dose in multiparous women and birthing people. (One complete Cycle).

Further Dinoprostone medication must be discussed with the Consultant/Senior Obstetrician.

Propess® (Dinoprostone)

Propess® is a pessary containing 10mg of Dinoprostone for release over 24 hours. Propess® should be considered as the first choice option for women and birthing people with a Bishop's score <4 (see further flowchart).

Administration: One pessary is inserted high into the posterior fornix. (One Cycle).

Propess® should be Removed

- After 24 hours
- When labour is established
- The membranes have ruptured and there are regular contractions >4 in 10
- If no regular contractions, Cervix less than 3cm dilated and CTG normal – **Propess® can stay in situ** until transfer to Labour room and Oxytocin drip arranged
- Significant vaginal bleeding
- There is evidence of uterine tachysystole, hypertonus or hyperstimulation
 - Tachysystole = ≥ 5 contractions in 10 minutes with normal CTG
 - Hypertonus = painful contraction lasting ≥ 90 seconds: normal CTG
 - Hyperstimulation = Tachysystole or hypertonus with abnormal CTG
- CTG suggests fetal hypoxia

- There is evidence of maternal systemic adverse effect such as severe nausea and vomiting

Artificial Rupture of the Membranes (ARM)

ARM should be considered as the first method of induction if the woman and birthing person's Bishops score is 7 or more.

If women and birthing people are suitable for ARM and oxytocin alone, an intravenous fluid chart and the oxytocin is to be prescribed and placed in the notes ready for admission.

Prior to ARM, abdominal and vaginal examination should be performed to confirm presentation and engagement. ARM should be discussed with the woman and birthing person and verbal consent obtained and this should be recorded in the notes.

When performing ARM a sterile technique and an Amnihook should be used to rupture the forewaters. Note in patients labour notes the cervical findings, indication, amount and colour of liquor and presence of meconium.

Cervical Ripening Balloon

Balloon can be recommended as the first line for IOL in specific circumstances i.e vaginal birth after caesarean, IUGR, grand multiparity, sensitivity to prostaglandin

Insertion of a double balloon catheter for induction of labour at term in pregnant women and birthing people aims to facilitate induction through causing dilation of the cervix when the cervix is unfavourable for induction. The double balloon is claimed to stimulate local prostaglandin release, which leads to cervical ripening, through the two balloons squeezing the cervix.

Please see link for full guidance: <http://www.nice.org.uk/guidance/ipg528> (opens in new tab).

Assessment

The components of the Bishops Score must be recorded in full (as below) using the sticker available on the Ward.

Modified Bishops Score (Max 10)	0	1	2
Dilatation (cm)	<1	1 – 2	3 – 4
Length of cervix (cm)	>4	2 – 4	1 – 2
Station (relative to ischial spines)	-3	-2	-1/0
Consistency	Firm	Average	Soft
Position	Posterior	Mid/Anterior	Anterior

Process of Induction of Labour

- Prostaglandins should be prescribed on the drug chart placed in patient's notes in preparation for ward admission (2 doses of Prostin E2® vaginal gel 1-2 mg or Propess® 10mg, see further in the flowchart)

- The rationale for induction of labour must be clearly discussed with women and birthing people and should include an appropriate individualised discussion around the risks and benefits. Women and birthing people should be counselled on the potential time for the induction process and women and birthing people should be advised to contact the maternity unit on the morning that their induction is booked, they should be informed that whether the induction will be able to go ahead will depend on the unit acuity
- IOL book to be completed in full
 - Name and address
 - Telephone number
 - Consultant
 - Gravida/para
 - Indication for IOL
 - Team
- Women and birthing people should be advised to ring for bed availability prior to admission. Admission time varies per local protocol. After admission to the ward, CTG with Dawes Redman criteria and baseline observations are commenced (BP, pulse, temperature, urinalysis, palpation).
- Assess CTG for 30 minutes with the view to proceed with IOL, perform vaginal examination and assess Bishop Score. Proceed with induction as per flow chart.
- After 30 minutes encourage mobilisation and await events.
- Intermittently auscultate the fetal heart according to clinical judgement.
- Dawes Redman criteria should not be used once the induction has commenced

Fetal Monitoring

Once regular contractions are reported fetal heart rate should be monitored with CTG for 30 minutes to assess fetal well-being.

Once in established labour, fetal heart to be monitored according to NICE intrapartum care guidelines

For women and birthing people who do not have any additional risk factor as referenced in the NICE intrapartum care guideline [2017] then intermittent auscultation can be used to assess the fetal well-being once labour is established without the use of oxytocin

Advise continuous cardiotocography if any of the following risk factors are present at initial assessment or arise during labour:

- Maternal pulse over 120 beats/minute on 2 occasions 30 minutes apart
- Temperature of 38°C or above on a single reading, or 37.5°C or above on 2 consecutive occasions 1 hour apart
- Suspected chorioamnionitis or sepsis

- Pain reported by the woman and birthing person that differs from the pain normally associated with contractions
- The presence of meconium
- Fresh vaginal bleeding that develops in labour
- Severe hypertension: a single reading of either systolic blood pressure of 160 mmHg or more or diastolic blood pressure of 110 mmHg or more, measured between contractions
- Hypertension: either systolic blood pressure of 140 mmHg or more or diastolic blood pressure of 90 mmHg or more on 2 consecutive readings taken 30 minutes apart, measured between contractions
- A reading of 2+ of protein on urinalysis and a single reading of either raised systolic
- blood pressure (140 mmHg or more) or raised diastolic blood pressure (90 mmHg or more)
- Confirmed delay in the first or second stage of labour (see recommendations)
- Contractions that last longer than 60 seconds (hypertonus), or more than 5 contractions in 10 minutes (tachysystole)
- Oxytocin use

Management of Uterine Hyperstimulation

- Tachysystole = ≥ 5 contractions in 10 minutes with normal CTG
 - Hypertonus = painful contraction lasting ≥ 90 seconds: normal CTG
 - Hyperstimulation = Tachysystole or hypertonus with abnormal CTG
1. Keep on CTG
 2. Place patient in left lateral position
 3. Remove Propess® – DO NOT perform ARM
 4. Inform on-call registrar and transfer to labour ward
 - IV Access (Take bloods for FBC / Group & Save)
 - IV 0.9% sodium chloride – start at a rate of 500ml/hr
 5. **If CTG is normal:**
 - Wait for 15-30 mins then reassess.
 - If tachysystole or hypertonus persisting, administer 250 micrograms SC **TERBUTALINE** (ie 0.5ml of 0.5mg/ml solution of Bricanyl®).
 6. **If CTG is NOT normal (suspicious or pathological):**
 - Administer 250 micrograms SC **TERBUTALINE** immediately, and involve the on call anaesthetist.

Unsuccessful IOL

Unsuccessful IOL is when labour is not established / or ARM is not possible after one cycle of IOL. If the IOL is unsuccessful, the woman and birthing person should have a full assessment by a senior obstetrician. This should include the indication for IOL, the method used, the maternal and fetal wellbeing, and the woman and birthing person's wishes.

The following options should be offered to the woman and birthing person:

Further Cycle of IOL

The timing and method of any further cycle should be determined on individual bases

Category 3 Caesarean Birth

The timing of the Caesarean should take into consideration the maternal and fetal wellbeing as well as the work load on the unit/neonatal unit.

Auditable Standards

- Number of women and birthing people having membrane sweeps prior to IOL
- Computerised CTG performed prior to commencement of induction
- Delays during the IOL procedure
- IOL Pathway Compliance

References

NICE [NG207] Inducing labour | Guidance | 2021 <https://www.nice.org.uk/guidance/ng207> (opens in new tab)

Bahtiyar MO et al Stillbirth at Term in Women and birthing people of Advanced Maternal Age in the United States: When Could the Antenatal Testing Be Initiated? *Amer J Perinatol* 2008;25(5): 301-304

Calder AA, MacKenzie IZ. Review of Propess - a controlled release dinoprostone (prostaglandin E₂) Pessary. *J Obstet Gynaecol.* 1997 Sep;17 suppl 2:S53-67.

Hilder L, Costeloe K, Thilaganathan B. Prolonged pregnancy: evaluating gestation-specific risks offetal and infant mortality. *BJOG: anInternational Journal of Obstetrics and Gynaecology* 1998;105(2):169–73.

Jacobsson B, Ladfors L, Milsom I. Advanced maternal age and adverse perinatal outcome. *Obstet Gynecol* 2004; 104:727–733.

Joseph KS, Allen AC, Dodds L, et al. The perinatal effects of delayed childbearing. *Obstet Gynecol* 2005; 105:1410–1418.

Mukhopadhyay M, Lim KJ, Fairlie FM. Is propess a better method of induction of labour in nulliparous women and birthing people? *J Obstet Gynaecol.* 2002 May;22(3):294-295.

NICE (2013) Evidence Update 44 Induction of Labour

NICE (2015) Insertion of a double balloon catheter for induction of labour in pregnant women and birthing people without previous caesarean section. NICE Interventional Guidance (IPG528)

NICE (2021) Inducing Labour (NG207)

NICE (2014) Intrapartum care for healthy women and birthing people and babies (NG190)

The Summary of Product Characteristics for Propess® 10mg vaginal delivery system, Ferring Pharmaceuticals Ltd. Date of revision: September 2021. Accessed via: www.medicines.org.uk/emc/product/135/smpc (opens in new tab)

The Summary of Product Characteristics for Prostin E2 Vaginal Gel 2mg, Pfizer Limited. Date of revision: July 2021. Accessed via: www.medicines.org.uk/emc/product/1090/smpc (opens in new tab)

Appendix 1 – IOL Sticker

Induction of Labour Decision			
Name:		Hospital number	
Indication	Gestation now	EDD	
	Gestation at induction		
	Date of induction		
Risk Factors			SCBU informed? Y N N/A
Potential Risks explained: More painful /hyperstimulation/may not be successful/Take longer/risks of Instrumental delivery, CS/others			
Discussion completed and documented (including procedure)		Consent obtained	
Information leaflet given		Prescription completed	
Consultant informed		Induction booked	
Woman informed that alternative options available		If IOL unsuccessful options discussed	
Pain relief options discussed		Plan for membrane sweeping clearly discussed and documented	

Appendix 2 – Bishop’s Score Sticker

Modified Bishops Score (Max 10)	0	1	2
Dilatation (cm) Length of cervix (cm)	<1 >4	3 – 2 4 – 4	3 – 4 1 – 2
Station (relative to ischial spines)	-3	-2	-1/0
Consistency	Firm	Average	Soft
Position	Posterior	Mid/Anterior	Anterior

Appendix 3 – Induction of Labour Information Leaflet (English)



GIG
CYMRU
NHS
WALES

Bwrdd Iechyd Prifysgol
Hywel Dda
University Health Board

Induction of Labour Information Leaflet

Your induction of labour has been booked for:

Date:

Time:

Please phone Dinefwr ward @ 0830 on the morning
of your induction to confirm your bed and the time to
come in.

Telephone no: 01267 283135




Induction of Labour (IOL)

There are many reasons why your labour may need to be induced which will be discussed with you by your midwife or doctor. It is important that you understand why you are being offered an induction of labour so if you are unsure please ask the person who is booking it for you to explain further. The aim of this leaflet is to help prepare you for what to expect during the induction of labour process.

What is Induction of Labour (IOL)?

IOL is the process used to start labour artificially. The aim is to help ripen your cervix to prepare it for labour. Your cervix is the bottom part of your uterus (neck of the womb) that opens to allow the baby to move from your uterus and down the birth canal to be born. During pregnancy your cervix is long, closed and firm. Towards the end of your pregnancy it softens, or ripens. As you go into labour it will soften more, get shorter and open up (dilate). It is not unusual for IOL to take up to 4-5 days although this is different for everyone. Please bring plenty of snacks and things to keep you occupied. You are welcome to bring your iPad or laptops but we do ask that you use headphones so you don't disturb the other people in your bay.



How is labour induced?

There are different methods we use to induce labour. Which method is most appropriate for you will depend upon an assessment of your cervix via a vaginal examination by your midwife. Before this examination your baby's heartbeat will be monitored for 30 minutes and again for 30 minutes afterwards. During the assessment we will give your cervix a score called the Bishop score which will help us choose the best option for your induction. The most appropriate option will then be discussed with you before continuing. You may require one or all of the following methods.


Method 1 = Propess (aka the teabag)

Propess is a vaginal pessary, which looks like a small tampon that is positioned to sit behind your cervix. The propess will slowly release the hormone prostaglandin over 24 hours to help ripen your cervix. There is a long string attached to the propess so it can easily be removed. It is important that you are careful not to pull or drag the string as this may cause the propess to come out. Please take care when wiping yourself after going to the toilet, after washing and when getting on and off the bed. If your propess does come out please inform the midwife straight away so it can be reinserted. After 24 hours you will be assessed again and may require a second propess which again will stay in for 24 hours.




Method 2 = Prostin gel

Prostin gel is inserted via an applicator similar to a syringe. The top of the applicator is aimed behind your cervix and the gel is released. After 6 hours you will be assessed again to see if you require another prostin gel.




Method 3 = Cervical Balloon

A cervical balloon is a mechanical way of inducing your labour as it does not involve any medications. This procedure is performed on labour ward by the doctors and involves a soft rubber tube being inserted through the cervix. Once in position the doctor will then inflate two balloons inside the tube, one above the cervix and one below the cervix. The balloon rubs against and stretches the cervix causing it to produce a hormone called prostaglandin. The prostaglandin causes the cervix to become shorter and soften (ripening). This prepares the cervix for labour and allows your midwife or doctor to break your waters. Once the balloon is in you will return to the ward overnight before going back to labour ward in the morning to have the balloon removed and your waters broken. You will usually be offered a balloon if there is a reason why you would benefit from not having the medications. For example if you have had a previous c/section to avoid putting any strain on your scar, or if you have had a reaction to the induction medications which has made you contract too much.



Method 4 = Artificial rupture of membranes (ARM)

This means breaking your waters. Once your cervix has thinned down and dilated to about 2-3cm we will be able to break your waters. This is done on labour ward by performing a vaginal examination and inserting a small plastic hook and putting pressure on the bag of membranes around the baby to make a hole. Once this is done we will usually allow you a couple of hours to mobilise in order to bring on contractions.



Method 5 = Oxytocin drip

This is a hormone drip which we will slowly increase every 30 minutes until we have 4 regular contractions every 10 minutes. This is done on labour ward and will involve having a cannula, usually in the back of your hand, so we can give you the hormone straight into your vein. At this point your baby's heartbeat will be continuously monitored. You can still remain mobile while you are on the monitor and the midwife looking after you can show you how to make full use of the birthing bed to adopt different positions.



What pain relief options are available?

Everyone experiences pain differently, so where most will find examinations mildly uncomfortable, some may find it painful. If this is the case for you please discuss this with your midwife so pain relief can be offered to you.

After the Propess/prostin is given you may feel some crampy 'period' like pains, backache, tightenings and some discomfort in you vagina. Walking around can help to ease your discomfort and can also help to get your labour established. Relaxation and breathing methods can be used, you can learn more about these at your antenatal classes or from your midwife. Adopting different positions can also help and we do have a number of birthing balls on the ward that you are free to use.

We do have a short supply of TENS machines available but these cannot be guaranteed. You can hire a TENS machine from some pharmacy outlets such as Boots as you may prefer to bring your own.

We recommend using water as a method of pain relief so would encourage you to use the bath or the showers.

If needed we can give you simple analgesia such as paracetamol.

If you require further pain relief you can discuss this with your midwife.



What are the risks of IOL?

- Induction of labour is a medical intervention which can affect your birth choices and experience.
- Induction may increase your chance of needing an assisted vaginal birth with forceps or ventouse.
- Induced labour may be more painful than spontaneous labour.
- Prostaglandins can cause vaginal soreness or irritation.
- Propess/prostin may cause your uterus may contract too frequently or for too long which may lead to baby becoming distressed. If this happens we will need to reverse the effects by removing the propess and/or giving you an injection to slow the rate of contractions.
- Induction is not always successful.
- Induction can take up to 5 days so you may have a long stay in hospital.

What if my induction doesn't work?

This doesn't happen very often but when it does the doctors will look at your individual case. Depending on your wishes you may be offered a day of rest before having some more of the induction medications or you will be offered a caesarean birth.

Can my birthing partner stay with me?

Unfortunately due to ongoing COVID restrictions your birthing partner will be unable to stay with you overnight during your induction. Our current visiting times for your birthing partner only is 10am - 8:30pm. We understand that this is a very anxious time for you and we are here to support you as much as we can. You are free to go off the ward to visit your birthing partner but we do ask that you limit the amount of times you are in and out to help reduce the spread of COVID. Once you go over to labour ward, either to have your waters broken or in established labour, your birthing partner will be allowed in to be with you. These restrictions are subject to change so please speak to your midwife about any current changes.

Is there a chance my induction will be delayed?

Unfortunately when the unit is very busy we may need to delay your induction so you need to be aware that your induction date is not guaranteed. We understand that this is very disappointing however the safety of you and your baby is most important and so sometimes this decision is the safest option due to the pressures on the unit. We ask you to phone the ward on the morning of your induction to check that we have a bed for you. We will then advise you on when to come in or whether there will be any delays. This does not happen very often but when it does the consultant on call will look at each individual case and prioritise who needs to come in first.

Do I have any other options?

You can choose to decline or delay your IOL. Please discuss this with your care provider so you can make an informed choice and an alternative plan of care can be made with you. If you have already started the induction process but decide you do not wish to continue your options will be discussed with you by the on call team.