

Gynae Datix trigger list

<u>Clinical</u>	<u>Organisational</u>
Unexpected death	Inappropriate/Lack of professional involvement
Loss of fetal remains	Equipment/facilities failure
Delayed or missed diagnosis	Patient identification issues
Damage to bowel/bladder/major blood vessel/uterus	Failure to follow local protocol
Bleeding more than expected >500 ml . If complicated case then more than expected of 1 ltr.	Staffing issues
Unplanned readmission within 30 days of surgery	Cancellation of elective procedures
Emergency re attendance in A&E or clinic requiring further treatment following colposcopy treatment	Clinics cancelled at less than 6 weeks
Drug Incident	Failure to escalate or request assistance
Unplanned return to theatre	Complications of anaesthesia
Unplanned HDU/ITU admission	Infection control/ cleanliness issues
Deviation from planned surgery	Potential service user complaint
Retained swab /instrument	
Haemorrhage requiring blood transfusion	
Abdominal wound complication- Wound dehiscence/burst abdomen	
Post op DVT /PE	
Failure to give anti-D	
Problems related to miscarriage/MTOPS/STOP/ERPC	
Omission of planned procedures (Violation of local protocol) (eg: failure to insert planned intrauterine contraceptive device after a hysteroscopy)	
Procedure performed without consent (ie: ovaries removed at hysterectomy)	

Author- Monika Vij

Version 1.2 20/01/2026

Review date 20/01/2029

Reference- RCOG clinical governance advice –Sep 2009