



GIG  
CYMRU  
NHS  
WALES

Bwrdd Iechyd Prifysgol  
Bae Abertawe  
Swansea Bay University  
Health Board

## **Departments of Uro-Gynaecology and Urology**

### **Managing recurrent urinary tract infection Advice for Patients**

Urine will never be completely sterile when you have a urostomy and this is not harmful. The aim is to keep symptom free and not to have sterile urine.

1. A simple strategy to stop an early infection is to increase your fluid intake
2. When you think you have a urine infection you should collect a mid stream sample using a red top bottle. Take this sample to your GP surgery, properly labeled, in order that it can be sent to the laboratory, and collect a new sample bottle to keep at home
3. Whilst waiting for the result, start the antibiotics that your last infection was sensitive to
4. Stop these antibiotics when your symptoms have settled. Sometimes this will be within 1 to 3 days
5. After 4 days your GP surgery will advise you of the result of your urine culture and whether you need a new prescription
6. The GP will advise you whether you need to take these tablets immediately or keep them available for your next infection
7. You can safely store antibiotics at home and they will remain useful for several years
8. Keep a bladder infection diary to record the dates of infection, record the results of samples, antibiotics and dose used, and the length of treatment
9. If you are female, over 45 years and menopausal, low dose vaginal oestrogen cream can protect against infection – ask your GP about this
10. Cranberry juice or tablets are helpful for some but not all patients – take advice from your health care professional

Contact Numbers for Advice:

Continence Service Morriston Hospital and Neath/Port Talbot Hospital

01792 703092

Continence Service Singleton Hospital  
Continence Service Community  
Urology Department Bridgend

01792 285384  
01639 683063  
01656 752470