





**All Patients with diabetes should have their blood glucose monitored a minimum of four times a day for the first 48 hours of admission, test pre-prandial and bedtime.  
Thereafter, monitor according to the capillary glucose monitoring guidance which is accessible on COIN.**

Date DD/MM/YY																					
Test time In 24hr																					
<b>BLOOD GLUCOSE</b> mmol per Litre	28																				
	26																				
	24																				
	22																				
	20																				
	18																				
	16																				
	14	REVIEW TREATMENT REGIME AND REFER TO DIABETES TEAM																			
	12																				
	10																				
	8																				
6																					
<b>IF CBG <math>\leq</math> 4mmol/L TREAT</b>	4																				
	2	TREAT HYPOGLYCAEMIA IMMEDIATELY AS DIRECTED BY GUIDANCE ON COIN																			
	1																				
CB Glucose																					
CB Ketones																					
Tester Print name Signature Role	AN EXAMPLE <i>A Example HCSW</i>																				

Date DD/MM/YY																					
Test time In 24hr																					
<b>BLOOD GLUCOSE</b> mmol per Litre	28																				
	26																				
	24																				
	22																				
	20																				
	18																				
	16																				
	14	REVIEW TREATMENT REGIME AND REFER TO DIABETES TEAM																			
	12																				
	10																				
	8																				
6																					
<b>IF CBG <math>\leq</math> 4mmol/L TREAT</b>	4																				
	2	TREAT HYPOGLYCAEMIA IMMEDIATELY AS DIRECTED BY GUIDANCE ON COIN																			
	1																				
CB Glucose																					
CB Ketones																					
Tester Print name Signature Role	AN EXAMPLE <i>A Example RGN</i>																				

SBUHB ThinkGlucose Non-Insulin Blood Glucose Monitoring Chart V2 2021