Antenatal mental health: an obstetric perspective

Sarah Myfanwy Pitts Consultant obstetrician & gynaecologist

Interests

None to declare



outline

Antenatal mental health & intergenerational effects



Change of practice?

Intergenerational effects of deprivation and poverty THE 1001 CRITICAL DAYS MANIFESTO IS CALLING FOR A REFOCUSING OF SUPPORT FOR A BABY'S FIRST 1001 DAYS.

IDENTIFYING NEED EARLY IS CRITICAL TO PREVENTING THE ABUSE AND NEGLECT OF BABIES AND IMPROVING THEIR EMOTIONAL WELLBEING.



A BABY'S DEVELOPMENT CAN BE DRAMATICALLY IMPROVED WITH EARLY AND EFFECTIVE SUPPORT FOR PARENTS.

PREGNANCY AND THE SECOND YEAR ARE A CRITICAL STAGE IN A CHILD'S DEVELOPMENT.

DAMAGE EARLY ON CAN CAUSE STRESS-RELATED CONDITIONS IN ADULT LIFE, SUCH AS HEART DISEASE OR SUBSTANCE ABUSE.

3

2

INFANTS AS YOUNG AS ONE CAN EXPERIENCE TRAUMA FROM WITNESSING DOMESTIC ABUSE.



ADAPTED FROM THE NSPCC'S 2011 ALL BABIES COUNT CAMPAIGN

HIGH LEVELS OF STRESS

IN EARLY CHILDHOOD CAN BE 'TOXIC' TO THE DEVELOPING BRAIN. OF SERIOUS CASE REVIEWS - INTO DEATHS OR SERIOUS ABUSE - INVOLVE A CHILD UNDER ONE.

36%

0

AT LEAST ONE OF THESE ISSUES APPEARS IN OVER 70% OF CASES WHERE A BABY HAS BEEN KILLED OR SERIOUSLY INJURED.



26% OF BABIES IN THE UK HAVE A PARENT AFFECTED BY DOMESTIC VIOLENCE, MENTAL HEALTH OR DRUG/ ALCOHOL PROBLEMS.

DOMESTIC VIOLENCE AFFECTS



MENTAL HEALTH PROBLEMS AFFECT 144,000 BABIES

PROBLEMS AFFECT 109,000 BABIES

Prenatal stress associated in child with increased

- Anxiety & depression
- Behavioural problems: ADHD, conduct disorder
- Impaired cognitive development, esp. language
- Sleep problems in infants
- Neonatal behaviour
- More difficult infant temperament
- Victimisation in school
- Schizophrenia
- ? Autism

... associated with

Reduced birth weight & GA Mixed handedness Altered finger print pattern Decreased telomere length Altered microbiome Asthma Altered immune function

ALSPAC

Avon Longitudinal Study of Parents and Children


Large prospective birth cohort
 - c. 14,000 pregnant woman
 - Rrecruited Bristol in 1990 - 1991

 Detailed information on children at 4, 7, 11 & 13 years old

Cohort with complete data n = 7,363

- Maternal postnatal anxiety & depression
- Paternal pre- and postnatal anxiety
- Parenting
- Maternal age & education
- Birth weight
- Smoking & alcohol
- Psychosocial factors: crowding (SES)

ALSPAC Predicted population prevalence of a probable mental health disorder in children born to high (top 15%: open bars and low prenatal anxiety (full bars) mothers. Based on SDQ scores. Results similar with prenantal depression





DON'T LISTEN TO "OLD WIVES' TALES"

No shock can mark an unborn baby. No horrible sight can deform him. BUT

WORRY, FEAR and ANGER may affect his mothers blood, which supplies his food. THEREFORE she should be CALM, HAPPY and SWEET-TEMPERED. Educational poster for expectant mothers

National Child Welfare Association

1919

the age old question

Nature

OR

Nurture

the new question

Nature

AND


Nurture


Epigenetics


Courtney Griffiths – 2014

https://www.youtube.com/watch?v=JTBg6hqeuTg

Histone modification







Modifiable by

MealsMindMuscle

The intra-uterine environment



Foetal programming

 Environment in the womb can alter the development of the fetus with a long lasting effect on the child


But it is not irreversible



cortisol

O'Donnell

- 2011
- Animation
- <u>http://www.beginbeforebirth.org/thescience/plac</u>
 <u>enta</u>



 Higher in utero exposure to cortisol is associated with lower cognitive function

 Sensitive early mothering can reverse the effects



Transgenerational inheritance

 When epigenetic `marks' laid down on sperm / eggs

Animal lab studies
 Dutch famine study


 You are what your grandmother ate

 Male smoking <11 y.o. ALSPAC

 1st & 2nd generation

Known costs of perinatal mental health problems per year's births in the UK, total: £8.1 billion


LSE, 2014 The cost of perinatal mental health problems





Costs v improvement The cost to the public sector of perinatal mental health problems is 5 times the cost of improving services.

Perinatal depression care (from Gavin, Meltzer-Brody, Glover, and Gaynes 2015)



Prescribing in the perinatal period



(un)safe in pregnancy / lactation

...and they lived happily ever after.....because of her anti-depressants.

THE END





Studies focus on risk & not benefit

resources

http://www.choiceandmedication.org /ncmh/

<u>http://www.uktis.org/</u>

What works with modern mums

- Facebook
- App store
- Netmums
- Twitter

 Online resources: <u>http://www.twoinmind.org/what-we-do</u>
 Dads

How women can help themselves

 Take time for themselves each day to relax

- Yoga
- Massage

Sitting down and listening to music

in your ANC?

Different for each mum
Discuss relationships & remember dads
Pregnancy: not just the medical / physical
Consider emotional transition

in your wider practice?

Pregnancy prevention? – Postnatal wards?

Pre-conceptual counselling? – EPAU?

thank you

Questions?